



King's Research Portal

DOI:

[10.1016/S2468-2667\(18\)30112-9](https://doi.org/10.1016/S2468-2667(18)30112-9)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gulliford, M., & Ravindrarajah, R. (2018). Frailty: From clinical syndrome to epidemiological construct? The Lancet Public Health. DOI: 10.1016/S2468-2667(18)30112-9

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Frailty: from clinical syndrome to epidemiological construct?



More people now live to advanced ages. The health and social care needs of the growing population with age-related frailty are becoming important public health and health policy concerns. Frailty has been described as "a distinctive health state related to the ageing process in which multiple body systems gradually lose their in-built reserves,"¹ rendering frail individuals susceptible to adverse outcomes such as falls, fractures, hospital admissions, and mortality. The notion of frailty has potential for risk stratification, but how can it be defined and recognised and what are the causes of frailty?

A qualitative study,² published earlier this year, found that physicians can generally recognise frailty when they see it in their patients but clinicians acknowledge uncertainty and lack of reliability in classifying their patients' frailty status in the absence of clear definitions and diagnostic criteria. Many researchers have tried to operationalise definitions of frailty, of which the most widely accepted is the frailty phenotype proposed by Fried and colleagues.³ This model is based on the co-occurrence of at least three of five apparently non-specific features including unintentional weight loss, self-reported exhaustion, weakness (low grip strength), slow walking speed, and low physical activity.³ Another approach to the classification of frailty draws on the cumulative deficit model leading to the calculation of a quantitative frailty index. The Frailty Index developed by Rockwood and colleagues,⁴ considers the number of possible deficits in an individual. Recent exemplars of this approach include the e-Frailty Index⁵ and the Hospital Frailty Risk Score.⁶ The construction of these indices shows considerable overlap with the concept of multiple morbidity.⁷ Frailty indices are generally strongly associated with mortality⁵ and this finding suggests that some patients with frailty might be approaching a stage of terminal decline.⁸

Two papers, one by Eric Brunner and colleagues⁹ and the other by Peter Hanlon and colleagues,¹⁰ in this issue of *The Lancet Public Health*, explore the epidemiology of frailty in the context of two well-known UK cohort studies, the Whitehall II and UK Biobank studies, respectively. Both reports assessed relatively young populations. The Whitehall II study assessed participants at a mean age of 69 years, with a minority of women, whereas the analysis of UK Biobank study included

participants aged 37–73 years. Both reports used Fried's phenotype to assess frailty, classifying participants as pre-frail if they showed one or two features, and frail if they showed three or more. In the Whitehall II study,⁹ fewer than 2% showed evidence of frailty under the age of 65, increasing to more than 10% at 75 years or older. There were substantial inequalities in the occurrence of frailty, which was more frequent in women, ethnic minority groups and those with low employment grade. Low employment grade at age 50 years was associated with 2.60 times higher odds of later frailty (95% CI 1.89–3.58). Participants who had long-term conditions or lifestyle risk factors for long-term conditions at the age of 50 years were more likely to develop frailty in later life, and these characteristics largely accounted for inequalities in frailty. These findings are consistent with those of a recent international study,¹¹ which explored inequalities in walking speed in old age and support the view that social inequality has a major negative effect on healthy ageing.¹²

In the UK Biobank study,¹⁰ 16538 (3%) of 493737 participants met the criteria for frailty, with more than a third of the sample (185360 [38%] of 493737) meeting the criteria for pre-frailty. Multiple morbidity was strongly associated with frailty in the UK Biobank data with participants with four or more long term conditions having 27 times higher odds of frailty than those with no long-term conditions (OR 27.1, 95% CI 25.3–29.1). Of all individual long-term conditions, frailty was most frequently observed in patients with multiple sclerosis and chronic fatigue syndrome. Obesity at midlife was associated with frailty in both studies, suggesting that targeting modifiable risk factors at midlife might reduce the occurrence of frailty at later ages.

These new analyses advance our understanding of the association between long-term conditions, multiple morbidity, and frailty. These studies show that long-term conditions and their risk factors are often antecedents of frailty; multiple morbidity and frailty often co-exist. The data also raise questions concerning our present epidemiological definitions of frailty. The high proportion of participants classified as pre-frail in the UK Biobank study could suggest that the definition lacks specificity; however, pre-frailty was associated with mortality in the UK Biobank data with hazard ratios for

Lancet Public Health 2018

Published Online

June 13, 2018

[http://dx.doi.org/10.1016/S2468-2667\(18\)30112-9](http://dx.doi.org/10.1016/S2468-2667(18)30112-9)

[http://dx.doi.org/10.1016/S2468-2667\(18\)30079-3](http://dx.doi.org/10.1016/S2468-2667(18)30079-3)

See Online/Articles

[http://dx.doi.org/10.1016/S2468-2667\(18\)30091-4](http://dx.doi.org/10.1016/S2468-2667(18)30091-4)

[http://dx.doi.org/10.1016/S2468-2667\(18\)30091-4](http://dx.doi.org/10.1016/S2468-2667(18)30091-4)

[http://dx.doi.org/10.1016/S2468-2667\(18\)30091-4](http://dx.doi.org/10.1016/S2468-2667(18)30091-4)

[http://dx.doi.org/10.1016/S2468-2667\(18\)30091-4](http://dx.doi.org/10.1016/S2468-2667(18)30091-4)

mortality at age 65–73 years compared with the non-frail group of 1.45 in men (95% CI 1.34–1.57) and 1.50 in women (1.34–1.68). Other recent studies have assessed the associations of low walking speed¹¹ or low grip strength¹³ with mortality without invoking the concept of frailty. There is a lack of consensus on the definition of frailty; the classification of individuals as frail depends on the theoretical construct incorporated in the frailty model,¹⁴ raising a possibility that there might be more than one kind of frailty. At present, the term frailty might be viewed as describing one phase of an ageing continuum and this continuum is perhaps not readily dichotomised.

**Martin Gulliford, Rathi Ravindrarajah*

School of Population Health and Environmental Sciences, King's College London, London, UK (MG); and Centre for Primary Care, Division of Population Health, Health Services Research and Primary Care, University of Manchester, Manchester, UK (RR)
martin.gulliford@kcl.ac.uk

We declare no competing interests.

Copyright © 2018 The Authors. Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

- 1 British Geriatrics Society. *What is frailty?* London: British Geriatrics Society, 2018 <http://www.bgs.org.uk/frailty-explained/resources/campaigns/fit-for-frailty/frailty-what-is-it> (accessed June 5, 2018).
- 2 Korenvain C, Famiyeh I-M, Dunn S, et al. Identifying frailty in primary care: a qualitative description of family physicians' gestalt impressions of their older adult patients. *BMC Family Practice* 2018; **19**: 61.

- 3 Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; **56**: M146–57.
- 4 Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci* 2007; **62**: 722–27.
- 5 Clegg A, Bates C, Young J, et al. Development and validation of an electronic frailty index using routine primary care electronic health record data. *Age Ageing* 2016; **45**: 353–60.
- 6 Gilbert T, Neuberger J, Kraindler J, et al. Development and validation of a Hospital Frailty Risk Score focusing on older people in acute care settings using electronic hospital records: an observational study. *Lancet* 2018; **391**: 1775–82.
- 7 Barnett K, Mercer SW, Norbury M, et al. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012; **380**: 37–43.
- 8 Diehr P, Williamson J, Burke GL, et al. The aging and dying processes and the health of older adults. *J Clin Epidemiol* 2002; **55**: 269–78.
- 9 Brunner EJ, Shipley MJ, Ahmadi-Abhari S, et al. Midlife contributors to socioeconomic differences in frailty during later life: a prospective cohort study. *Lancet Public Health* 2018; published online June 13. [http://dx.doi.org/10.1016/S2468-2667\(18\)30079-3](http://dx.doi.org/10.1016/S2468-2667(18)30079-3).
- 10 Hanlon P, Nicholl BJ, Jani BD, et al. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality: a prospective analysis of 493 737 UK Biobank participants. *Lancet Public Health* 2018; published online June 13. [http://dx.doi.org/10.1016/S2468-2667\(18\)30091-4](http://dx.doi.org/10.1016/S2468-2667(18)30091-4).
- 11 Stringhini S, Carmeli C, Jokela M, et al. Socioeconomic status, non-communicable disease risk factors, and walking speed in older adults: multi-cohort population based study. *BMJ* 2018; **360**: k1046.
- 12 Cooper R. Socioeconomic adversity—an important barrier to healthy aging. *BMJ* 2018; **360**: k1288.
- 13 Celis-Morales CA, Welsh P, Lyall DM, et al. Associations of grip strength with cardiovascular, respiratory, and cancer outcomes and all cause mortality: prospective cohort study of half a million UK Biobank participants. *BMJ* 2018; **361**: k1651.
- 14 Mitnitski A, Fallah N, Rockwood MR, et al. Transitions in cognitive status in relation to frailty in older adults: a comparison of three frailty measures. *J Nutrition, Health & Aging* 2011; **15**: 863–67.